Author Index

Abidi, S. L.
Reversed-phase thin-layer chromatography of homologues of antimycin A and related derivatives 453
Aerts, M. M. L., see Keukens, H. J. 149
Ahsan, T.
——, Colenutt, B. A. and Sing, K. S. W.
Application of gas chromatography for the study of precipitated calcium carbonate 416
Arm, H., see Däppen, R. 39
Ayyangar, N. R.
—— and Bhide, S. R.
Separation of 1-naphthylamine from the five known impurities and sub-ppm level detection and quantitation of 2-naphthylamine by normal-phase high-performance liquid chromatography 201
Bartha, A.
——, Billiet, H. A. H. and De Galan, L.
Practical use of an optimization strategy in reversed-phase high-performance liquid chromatography for the separation of a limited subset of components in a reaction mixture 225
Beek, T. A. van, see Van Beek, T. A. 375
Benassi, C. A.
——, Semenzato, A. and Bettero, A.
High-performance liquid chromatographic determination of free formaldehyde in cosmetics 387
Bermejo, J., see Díez, M. A. 93
Berridge, J. C., see Wright, A. G. 27
Bettero, A., see Benassi, C. A. 387
Bhide, S. R., see Ayyangar, N. R. 201
Billiet, H. A. H., see Bartha, A. 225
Blaakmeer, A., see Van Beek, T. A. 375
Blanco, C. G., see Díez, M. A. 93
Boutar, M., see Lebrun, M. H. 307
Buckley, J. J.
—— and Wetlaufer, D. B.
Use of the surfactant 3-(3-cholamidopropyl)-dimethylammoniopropane sulfonate in hydrophobic interaction chromatography of proteins 61
Buszewski, B.
——, Sieńko, D. and Suprynowicz, Z.
Effect of the structure and density of chemically bonded C_{18} phase on the recovery of tryptophan and its metabolites from human urine 73
Caude, M., see Jadaud, P. 333
——, see Morin, P. 125
Cavazza, M., see Zandomeneghi, M. 289
Cavazzutti, G., see Gagliardi, L. 428
Chrétien, J. R., see Walczak, B. 237
Colenutt, B. A., see Ahsan, T. 416
Compadre, C. M., see Fullas, F. 213
Däppen, R.
——, Meyer, V. R. and Arm, H.
New insights into the recognition mechanisms of chiral "brush-type" liquid chromatographic stationary phases 39
De Galan, L., see Bartha, A. 225
Díez, M. A.
——, Fañanas, F. J., Blanco, C. G. and Bermejo, J.
Separation and characterization of methylcyclopentadiene dimers by gas chromatography and molecular spectroscopy 93
Dohnal, J., see Jelínek, I. 139
Douse, J. M. F.
Dynamic headspace method for the improved clean-up of gunshot residues prior to the detection of nitroglycerine by capillary gas chromatography with thermal energy analysis detection 178
Dreux, M., see Walczak, B. 237
Drozd, J., see Vejrosta, J. 394
Duteurtre, X., see Jadaud, P. 333
Ensminger, W. D., see Stetson, P. L. 163
Fañanas, F. J., see Díez, M. A. 93
Felix, G., see Walczak, B. 237
Fell, A. F., see Wright, A. G. 27
Fullas, F.
——, Kim, J., Compadre, C. M. and Kinghorn, A. D.
Separation of natural product sweetening agents using overpressured layer chromatography 213
Gagliardi, L.
——, Cavazzutti, G., Montanarella, L. and Tonelli, D.
Determination of sun-screen agents in cosmetic products by reversed-phase high-performance liquid chromatography. Part II. 428
Galan, L., de, see Bartha, A. 225
Games, D. E., see Ramsey, E. D. 353
Gaudemer, A., see Lebrun, M. H. 307
Gaudemer, F., see Lebrun, M. H. 307

Gazdag, M.
—— and Szepesi, G.
Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. IV. Selection of most applicable separation system 279
——, see Szepesi, G. 265
Gendron, A., see Lamarre, C. 448
Graebe, J. E., see Odén, P. C. 195
Hadžija, O.
——, Iskrić, S. and Tonković, M.
Mobility of some carboxy- and hydroxybenzene derivatives on thin layers of plain and iron(III)-impregnated silica gel 220
Hancock, D. O.
—— and Synovec, R. E.
Microbore liquid chromatography and refractive index gradient detection of low-nanogram and low-ppm quantities of carbohydrates 83
Hasegawa, K., see Hayakawa, K. 343
Hayakawa, K.
——, Hasegawa, K., Imaizumi, N., Wong, O. S. and Miyazaki, M.
Determination of amphetamine-related compounds by high-performance liquid chromatography with chemiluminescence and fluorescence detections 343
Heindorf, M. A.
—— and McGuffin, V. L.
Limitations of 2-diphenylacetyl-1,3-indandione-1-hydrazone as a precolumn fluorescence derivatization reagent 186
Henoux, J., see Jadaud, P. 333
Imaizumi, N., see Hayakawa, K. 343
Iskrić, S., see Hadžija, O. 220
Jadaud, P.
——, Caude, M., Rosset, R., Duteurtre, X. and Henoux, J.
Determination of some polynuclear aromatic hydrocarbons in new, used and regenerated oils 333
Jelínek, I.
——, Dohnal, J., Snopek, J. and Smolková-Keulemansová, E.
Use of cyclodextrins in isotachophoresis. VII. Resolution of structurally related and chiral phenothiazines 139
Joelsson, M., see Johansson, G. 49
Johansson, G.
—— and Joelsson, M.
Partition of hydrophobic compounds between two liquid phases of similar hydrophobicity 49
Kawai, S., see Takayama, Y. 172
Keukens, H. J.
—— and Aerts, M. M. L.
Determination of residues of carazolol and a number of tranquillizers in swine kidney by high-performance liquid chromatography with ultraviolet and fluorescence detection 149
Kim, J., see Fullas, F. 213
Kinghorn, A. D., see Fullas, F. 213
Kleinjans, J. C. S., see Verhagen, H. 438
Koizumi, K.
——, Kubota, Y., Tanimoto, T. and Okada, Y.
High-performance anion-exchange chromatography of homogeneous D-gluco-oligosaccharides and -polysaccharides (polymerization degree $\geqslant 50$) with pulsed amperometric detection 365
Kubota, Y., see Koizumi, K. 365
Kysilka, R.
—— and Wurst, M.
High-performance liquid chromatographic determination of some psychotropic indole derivatives 434
Lafosse, M., see Walczak, B. 237
Lamarre, C.
—— and Gendron, A.
Analyse quantitative par chromatographie liquide haute performance du di-*tert.*-butyl-2,5-*para*-crésol dans les huiles de transformateurs 448
Lambert, W. J.
—— and Wright, L. A.
Prediction of alkane–water partition coefficients using a C_{18} derivatized polystyrene–divinylbenzene stationary phase 400
Lebrun, M. H.
——, Gaudemer, F., Boutar, M., Nicolas, L. and Gaudemer, A.
Ion-pair, anion-exchange and ligand-exchange high-performance liquid chromatography of tenuazonic acid and 3-acetyl 5-substituted pyrrolidine-2,4-diones 307
Lee, H. K., see Ong, C. P. 405
Lesser, J. H., see Shustina, R. 208
Li, S. F. Y., see Ong, C. P. 405
McGuffin, V. L., see Heindorf, M. A. 186
Meyer, V. R., see Däppen, R. 39
Mihályfi, K., see Szepesi, G. 265
Mikešová, M., see Vejrosta, J. 394
Miller, S. L., see Smith, R. M. 297
Miyano, S., see Yamashita, J. 411
Miyazaki, M., see Hayakawa, K. 343
Montanarella, L., see Gagliardi, L. 428

Morin, P.
——, Pichard, H., Richard, H., Caude, M. and Rosset, R.
Supercritical fluid chromatography of sesquiterpene hydrocarbons on silica packed columns with on-line Fourier transform infrared detection 125
Morin-Allory, L., see Walczak, B. 237
Nicolas, L., see Lebrun, M. H. 307
Odén, P. C.
——, Schwenen, L. and Graebe, J. E.
Separation of gibberellins by normal-phase high-performance liquid chromatography 195
Ôi, S., see Yamashita, J. 411
Okada, Y., see Koizumi, K. 365
Ong, C. P.
——, Lee, H. K. and Li, S. F. Y.
Optimization of mobile phase composition for high-performance liquid chromatographic analysis of eleven priority substituted phenols 405
Peltonen, K.
Sampling and determination of carbon disulphide in air by gas chromatography with electron-capture detection 422
Perkins, J. R., see Ramsey, E. D. 353
Pichard, H., see Morin, P. 125
Rabuor, J. O., see Smith, R. M. 117
Ramsey, E. D.
——, Perkins, J. R., Games, D. E. and Startin, J. R.
Analysis of drug residues in tissue by combined supercritical-fluid extraction–supercritical-fluid chromatography–mass spectrometry–mass spectrometry 353
Richard, H., see Morin, P. 125
Rivas, J. D. L.
Reversed-phase high-performance liquid chromatographic separation of lutein and lutein fatty acid esters from marigold flower petal powder 442
Rosset, R., see Jadaud, P. 333
——, see Morin, P. 125
Satoh, H., see Yamashita, J. 411
Schwenen, L., see Odén, P. C. 195
Semenzato, A., see Benassi, C. A. 387
Seto, Y.
—— and Shinohara, T.
Size fractionation of oligosaccharides by liquid chromatography on a cation-exchange column 323
Shinohara, T., see Seto, Y. 323
Shukla, U. A., see Stetson, P. L. 163
Shustina, R.
—— and Lesser, J. H.
Liquid chromatographic determination of hydrazine, carbohydrazide and thiocarbohydrazide in aqueous solutions 208
Sieńko, D., see Buszewski, B. 73
Sing, K. S. W., see Ahsan, T. 416
Smith, R. M.
—— and Miller, S. L.
Comparison of the selectivity of cyano-bonded silica stationary phases in reversed-phase liquid chromatography 297
——, and Rabuor, J. O.
Separation of basic drugs by high-performance liquid chromatography on a silica column using a methanol–ethylenediamine buffer 117
Smolková-Keulemansová, E., see Jelínek, I. 139
Snopek, J., see Jelínek, I. 139
Startin, J. R., see Ramsey, E. D. 353
Stetson, P. L.
——, Shukla, U. A. and Ensminger, W. D.
Stability of trimetrexate, a new non-classical antifolate, in infusion solutions 163
Suprynowicz, Z., see Buszewski, B. 73
Sütfeld, R.
Preparative liquid chromatography with analytical separation quality. Interval injection/displacement reversed-phase high-performance liquid chromatography 103
Suzuki, T., see Yamashita, J. 411
Svoboda, V.
Ideal mixed cells model of multi-component preparative liquid chromatography 1
Synovec, R. E., see Hancock, D. O. 83
Szepesi, G.
——, Gazdag, M. and Mihályfi, K.
Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. III. Method validation 265
——, see Gazdag, M. 279
Takai, N., see Yamashita, J. 411
Takayama, Y.
——, Takeichi, T. and Kawai, S.
Application of deactivated metal capillaries to the analysis of solvents in varnish 172
Takeichi, T., see Takayama, Y. 172
Tanimoto, T., see Koizumi, K. 365
Tonelli, D., see Gagliardi, L. 428
Tonković, M., see Hadžija, O. 220
Yamashita, J.
——, Satoh, H., Ôi, S., Suzuki, T., Miyano, S. and Takai, N.
High-performance liquid chromatographic separation of enantiomers on axially chiral binaphthalene dicarboxylic acid–chiral phenylethylamine bonded to silica gel 411
Van Beek, T. A.
—— and Blaakmeer, A.
Determination of limonin in grapefruit juice and other citrus juices by high-performance liquid chromatography 375

Vejrosta, J.
——, Mikešová, M. and Drozd, J.
Sorption of *n*-alkanes on Tenax 394
Verhagen, H.
—— and Kleinjans, J. C. S.
Rapid determination of isomer ratios of butylated hydroxyanisole by high-performance liquid chromatography 438
Voelkel, A.
Properties of topological indices as structural parameters for compounds containing oligooxyethylene chain(s) 251
Walczak, B.
——, Dreux, M., Chrétien, J. R., Morin-Allory, L., Lafosse, M. and Felix, G.
Factor analysis and experiment design in high-performance liquid chromatography. VIII. Charge-transfer *versus* standard stationary phases in normal-phase high-performance liquid chromatography 237
Wetlaufer, D. B., see Buckley, J. J. 61
Wong, O. S., see Hayakawa, K. 343
Wright, A. G.
——, Fell, A. F. and Berridge, J. C.
Development and optimisation of a high-performance liquid chromatographic assay for tioconazole and its potential impurities. I. Selection of separation conditions 27
Wright, L. A., see Lambert, W. J. 400
Wurst, M., see Kysilka, R. 434
Zandomeneghi, M.
—— and Cavazza, M.
Chiral recognition by biological macromolecules. Partial resolution of racemic enones by albumin 289

Erratum

J. Chromatogr., 456 (1988) 444
Page 444, 3rd paragraph, 1st line, "0.01 *M*" should read "0.1 *M*".